

**PRESBYOPIA TREATMENT BY LENS ALTERATION**

**BACKGROUND OF THE INVENTION**

**Field of the Invention**

5           The present invention relates to a method and  
device for reversing and treating presbyopia.

**Background of the Invention**

Presbyopia affects virtually every person over the  
age of 44. According to Jobson Optical Database, 93% of  
10 people 45 and over are presbyopic. Presbyopia entails the  
progressive loss of amplitude of accommodation that  
occurs with aging. Adler's Physiology of the Eye, which  
is incorporated herein by reference, discloses that the  
human accommodative amplitude declines with age such  
15 that accommodation is substantially eliminated by the  
age of 50 to 55. Accommodative ability, as defined by  
U.S. Patent No. 5,459,133 to Neufeld and incorporated in  
its entirety herein by reference, is the capacity of the  
eye to focus for near vision by changing the shape of  
20 the lens to become more convex.

The ocular tissues involved in the accommodative  
response include the lens, the zonules, the lens  
capsule, and the ciliary muscle. Of these, the lens is

the central tissue. These structures function together to enable the eye to focus on close objects by changing the shape of the lens. The lens is centrally suspended between the anterior and posterior chambers behind the pupillary opening of the iris. The lens is supported by a radially oriented array of zonules which extend from the lateral edges of the lens to the inner border of the circumferential ciliary muscle. The ciliary muscle is attached to the scleral coat of the eye. When the eye is at rest, it is focused for distance and the lens is in a somewhat flattened or less convex position. This shape is due to the tension that is exerted on the lens' periphery by the zonules. The zonules pull the edges of the lens toward the ciliary body.

During accommodation, the shape of the lens becomes more convex through contraction of the ciliary muscle which allows the ciliary attachment of the zonules to move toward the lens, reducing the tension in the zonules. This reduction in tension allows the central region of the lens to increase in convexity, thereby enabling near objects to be imaged on the retina. The processes involving the coordinated effort of the lens, zonules, ciliary body, medial rectus muscles and iris,

among others, that leads to the ability of the eyes to clearly focus near on the retina is the accommodative process.

Several theories have been advanced to explain the  
5 loss of accommodation with age. These theories include the hardening of the lens with age; loss of strength in the ciliary muscle; and, the loss of elasticity of the lens capsule. As for the loss of strength of the ciliary muscle, it is noted that although there are age-  
10 related morphological changes that occur, there is little evidence of diminishing strength of the ciliary muscle. In fact, under the influence of pilocarpine, the ciliary muscle will vigorously contract even in presbyopic eyes.

15 As for changes in the lens capsule, it has been postulated that reduction in the elasticity of the capsule is, in fact, a contributing factor in presbyopia. Moreover, it has been found that Young's modules of elasticity for the lens capsule decreases by  
20 nearly 50% from youth to age 60, while accommodation decreases by 98%. Consequently, the principal cause of presbyopia is now considered to be "lenticular sclerosis" or the hardening of the lens. This hardening

of the lens may be due to an alteration of the structural proteins or an increased adhesion between the lens fibers.

A cataract is a condition in which the lens becomes  
5 less clear. The study of cataracts lends insight into lens and capsular changes. The usual senile cataract is relatively discus-shaped when removed from the eye, its shape being dictated by the firm lens substance. The liquefied hypermature cataract is globular when  
10 extracted, rounded up by the elastic lens capsule. This is indirect evidence that it may be possible to reverse the lenticular changes associated with presbyopia, and that the lens capsule is still sufficiently elastic.

Other theories advanced to explain presbyopia  
15 involve the role of lens growth through life and the loss of tension on the lens capsule. These theories, however, have not been supported by clinical observations.

At the present time, common treatments for  
20 presbyopia include reading glasses, bifocal glasses, or mono-vision contact lenses. All of these solutions necessitate the use of an appliance creating additional

shortcomings.

Alternative theories for treating presbyopia include scleral expansion and corneal reshaping. The efficacy of such techniques is not well-established and, importantly, these techniques do not attempt to reverse what the inventors of the subject application believe to be a substantial causation, as explained more fully below, in the loss of the accommodative amplitude of the lens typically associated with the normal aging process. Thus, whereas the present invention as explained further below, is directed to a method of reversing or treating presbyopia resulting in underlying changes in the structures and/or interactions of molecules comprising those components of the eye associated with the accommodative process, most notably the lens and/or lens capsule, scleral expansion and corneal reshaping involve macroscopic changes in the morphology of the lens and cornea. Thus, the present invention provides a novel molecular approach to reversing presbyopia by restoring the accommodative amplitude of the lens, and in another preferred embodiment, to reversing presbyopia by such novel approach while also reducing the tendency for the lens

to lose its thus restored accommodative amplitude.

Finally, the use of the excimer laser for the purposes of corneal reshaping to produce a multifocal refracting surface has been disclosed in Patent No.

5 5,395,356. While this method seems promising, it still requires structural changes to compensate for aging.

#### **SUMMARY OF THE INVENTION**

In its broadest sense, the present invention is directed to increasing the accommodative amplitude of  
10 the lens and thus to a method for reversing and/or treating presbyopia. In one embodiment, the present invention is directed to a method for reversing and/or treating presbyopia by breaking disulfide bonds in molecules comprising the structures of the eye, most  
15 notably the lens and the lens capsule, which disulfide bonds are believed to be a substantial factor in the progressive loss of accommodative amplitude. In another embodiment, the breaking of the disulfide bonds is accompanied by chemical modification of the free sulfur  
20 atoms formed upon breaking of the disulfide bonds, such chemical modification rendering the sulfur atoms less likely to form new disulfide bonds. This method thus comprises a method for preventing the recurrence of

presbyopia by reducing the availability of new disulfide bonds to be formed. Particularly, this invention effects a change in the accommodative amplitude of the human lens by: (1) using various reducing agents that cause a  
5 change in the accommodative abilities of the human lens, and/or (2) the use of external energy to affect a change in the accommodative abilities of the human lens. It is believed that by breaking bonds, such as disulfides, that adhere lens fibers together and cause a hardening  
10 of the lens cortex, the present invention increases the elasticity and the distensibility of the lens cortex and/or the lens capsule.

Presbyopia, or the loss of the accommodative amplitude of the lens, has often advanced in a typical  
15 person age 45 or older to the point where some type of corrective lens in the form of reading glasses or other treatment is required. It is to be understood that loss of accommodative amplitude can occur in persons much younger or older than the age of 45, thus the present  
20 invention is not to be construed as limited to the treatment of presbyopia in a person of any particular age. The present invention is most useful in a person whose accommodative amplitude has lessened to a point

where restoration thereof to some degree is desirable.

The accommodative amplitude of the lens is measured in diopters. The lens of a person who does not suffer from presbyopia (i.e. a person whose lens accommodates normally), will typically have an accommodative amplitude of about 2.5 diopters or greater. The terms "reversing presbyopia" or "treating presbyopia" as used in herein mean increasing the accommodative amplitude of the lens. The present invention is thus directed to a method for reversing presbyopia or increasing the accommodative amplitude of the lens of an individual. In a preferred embodiment of the present invention, the method of reversing presbyopia will result in an increase in the accommodative amplitude at least about 0.5 diopters. In a more preferred embodiment of the present invention, the method of reversing presbyopia will result in an increase in the accommodative amplitude of at least about 2.0 diopters.

In an even more preferred embodiment, the method of reversing presbyopia of the present invention will result in an increase in the accommodative amplitude by at least about 5 diopters. In a most preferred embodiment of the present invention, the method of



reversing presbyopia of the present invention will  
result in an increase of the accommodative amplitude of  
the lens to restoration thereof to that of a lens with a  
normal accommodative amplitude of 2.5 diopters or  
5 greater would result. While it is obviously most  
beneficial to restore the accommodative amplitude of the  
lens of each patient to a normal accommodative  
amplitude, it is to be understood that lesser degrees of  
restoration are also beneficial and in some cases due,  
10 for example, to a severe reduction in the accommodative  
amplitude (i.e. advanced presbyopia) it may not be  
possible to obtain full restoration to normal  
accommodative amplitude.

#### **DETAILED DESCRIPTION**

15 As stated, inelasticity of the lens, or hardening  
thereof, is believed to be a contributing cause of  
presbyopia. The hardening of the lens may be due to an  
alteration of the structural proteins or an increased  
adhesion between the lens fibers. In one embodiment,  
20 the present invention is directed to treating presbyopia  
by altering the molecular and/or cellular bonds between  
the cortical lens fibers so as to free their movement  
with respect to each other. The increased elasticity of

the lens apparatus can restore lost amplitude of accommodation. It is believed that disulfide bonds in the molecules comprising the structures of the eye responsible for proper accommodation are a substantial  
5 factor in the hardening of the lens and the concomitant loss of accommodative amplitude.

Thus, in one embodiment of the invention, a two-step process involves breaking the disulfide bond, and then protonating the newly-formed sulfur atom with a  
10 reducing agent such as glutathione to impart a hydrogen atom thereto. The steps can be performed simultaneously or consecutively. In either case, the reducing agent should be present at the time the disulfide bond is broken in order to eliminate reformation of disulfide.  
15 That is, the reducing agent introduces and bonds a moiety onto the free sulfur atom after breaking the disulfide bond such that the likelihood of reformation of another disulfide bond is prevented or at least reduced. While the reducing agent may introduce a  
20 hydrogen atom onto the free sulfur atom, thus forming a sulfahydryl (-SH group), the resultant -SH groups can again be oxidized to form a new disulfide bond. Thus, it is advantageous to introduce a group into the free

sulfur atom, such a -CH<sub>3</sub> or other moiety, that reduces the tendency of new disulfide bond formation. This method can result in a substantial prevention of the reoccurrence of presbyopia.

5           As stated, it is believed that the disulfide bonds form between the lens fibers and substantially reduce the lens fibers' ability to easily move relative to each other and thus the ability of the lens to accommodate properly. While not wishing to be bound by any  
10 particular theory, the bonds may form by way of absorption of light energy, which causes the sulfahydryl bonds on the lens proteins to oxygenate removing a hydrogen atom from two adjacent -SH groups and creating water and a disulfide bond. Reducing the disulfide  
15 bonds requires hydrogen donors such as glutathione or other molecules.

The total refractive power of the lens is greater than what would be expected based on the curvature and the index of refraction. As stated, contraction of the  
20 ciliary muscle causes the ciliary body to move forward and towards the equator of the lens. This causes the zonules to relax their tension on the lens capsule, which allows the central lens to assume a more spherical

shape. During accommodation, the main change is in the more central radius of curvature of the anterior lens surface, which is 12mm in the unaccommodative state and can be 3mm centrally during accommodation. Both the  
5 peripheral anterior and the posterior lens surfaces change very little in curvature during accommodation. The axial thickness increases while the diameter decreases. The central anterior lens capsule is thinner than the rest of the anterior capsule. This may explain  
10 why the lens bulges more centrally during accommodation. The thinnest portion of the capsule is the posterior capsule, which has a curvature greater than the anterior capsule in the unaccommodative state.



The protein content of the lens, almost 33% by  
15 weight, is higher than any other organ in the body. There are many chemical compounds of special interest in the lens. For example, glutathione is found in high concentration in the lens cortex even though there is very little in the aqueous. Thus, the lens has a great  
20 affinity for glutathione and actively absorbs, transports and synthesizes glutathione.

Approximately 93% of intralenticular glutathione is in the reduced form. Glutathione may be involved with

maintaining the lens proteins, the sulfahydryl groups  
(-SH), in their reduced states. That is, after the  
disulfide bond is broken and the sulfur atoms are made  
available, glutathione can impart a hydrogen atom to  
5 form the sulfahydryl group thereby preventing or  
minimizing the reformation of a disulfide bond. In  
addition, ascorbic acid can also be found in very high  
concentrations in the lens. It is actively transported  
out of the aqueous and is at concentrations 15 times  
10 that found in the bloodstream. Both inositol and  
taurine are found at high concentrations in the lens for  
which the reason is not known.

According to one embodiment of the invention , the  
increase in the accommodative amplitude is accomplished  
15 by treatment of the outer lens region (the cortex) with  
radiation, heat, chemical, enzyme, gene therapy,  
nutrients, other energy source, and/or any combination  
of any of the above sufficient to break the disulfide  
bonds believed responsible for the inelasticity of the  
20 lens. Chemicals are useful to reduce disulfide bonds  
that are believed to anchor lens fibers hence preventing  
their free movement and elasticity. By making the  
anterior cortex more elastic, viscosity is lowered and

10


$$\begin{array}{c} \text{R}_1 \\ \diagdown \\ \text{N} \\ \diagup \\ \text{R}_2 \end{array} - \text{C}(=\text{S}) - \text{S} - \text{S} - \text{C}(=\text{S}) - \begin{array}{c} \text{R}_3 \\ \diagup \\ \text{N} \\ \diagdown \\ \text{R}_4 \end{array}$$


wherein  $R_1$ ,  $R_2$ ,  $R_3$  and  $R_4$  are independently a  
straight or branched lower alkyl which may be  
substituted, e.g., by hydroxyl, lower alkoxy or lower  
alkyl carbonyloxy, their derivatives or a  
5 pharmaceutically acceptable salt thereof. Preferred  
exemplary reducing agents include diethyl  
dithiocarbamate, 1-methyl-1H-tetrazol-5-yl-thiol and 1-  
(2-hydroxyethyl)-1H-tetrazol-5-yl-thiol or and  
pharmaceutically acceptable salts thereof. Other useful  
10 compounds can be found in U.S. Patent No. 5,874,455  
which is hereby incorporated in its entirety by  
reference. The above-listed chemicals are merely  
exemplary and other reducing agents which behave  
similarly by breaking the disulfide bond are included  
15 within the scope of this invention.

The chemical reducing agents can be used alone or  
in conjunction with a catalyst such as an enzyme.  
Enzymes and other nutrients suitable for causing or  
facilitating reduction include, for example,  
20 aldoreductase, glyoxylase, glutathione S-transferase,  
thiol reductase, tyrosine reductase or any compatible  
reductase. Again, it should be noted that the above-  
listed enzymes are exemplary and not an exhaustive list.

The enzymes can be naturally present in the eye, or can be added to the eye together with or separate from the chemical reducing agent or energetic means disclosed herein. As such, other chemically and biologically comparable enzymes which help break disulfide bonds or behave similarly should be considered as within the scope of the present invention.

In one embodiment of the invention, the reduction of disulfide groups of the lens proteins to sulfahydryl groups is accomplished by delivering to the lens a compound such as glutathione, thiols, or others in sufficient quantities to reduce the disulfide bonds and other molecular and cellular adhesions. Other enzymes or chemicals that affect a methylation on the free sulfur atom include for example, methyl-methane tiosulfonate, methyl glutathione, S-methyl glutathione, S-transferase and other biologically compatible methylating agent. Use of emulsions such as nanocapsules, albumin microspheres, carrier molecules such as inositol, taurine or other biologically suitable means for delivering the reducing agent to the lens is an integral part of this invention. The chemical reducing agent will typically be delivered in the form



of a solution or suspension in an ophthalmically acceptable carrier. In some cases, the application of external energy to affect or catalyze the reduction of the disulfide bonds as well as the disruption of other  
5 bonds and adhesions, may be beneficial. The application of external energy alone can be used to break the disulfide bonds. Externally applied energy can have any form, by way of example only, any of laser, ultrasound, heat, ionizing, light, magnetic, microwave, sound,  
10 electrical, or other not specifically mentioned, can be used alone or in combination with the reducing agents to affect the treatment of presbyopia, or a combination of any of these types of energy.

In a similar manner, agents can be delivered to the  
15 lens capsule which bind or interact with the capsule to affect greater elasticity or distensibility. Such agents either cause the capsule to shrink in surface area or increase the tension of the lens capsule on the peripheral anterior or posterior of the lens.  
20 Externally applied energy can have any form, by way of example only, any of laser, ultrasound, heat, ionizing, light, magnetic, microwave, sound, electrical, or other not specifically mentioned can be used alone or in

combination with the reducing agents to affect the treatment of presbyopia or a combination of any of these types of energy.

In another embodiment of the invention,  
5 externally applied energy can be used as a catalyst to induce or increase the rate of the reduction reaction. Thus, by applying external energy, the peripheral portion of the capsule is preferentially affected, leaving the central 4mm zone of accommodation  
10 unaffected. This allows the lens to assume a more accommodative state. The externally applied energy can also be applied alone to promote the reduction reaction and the cellular changes that ultimately affect the lens' cortex.

15 As examples, lasers useful in the present invention include: excimer, argon ion, krypton ion, carbon dioxide, helium-neon, helium-cadmium, xenon, nitrous oxide, iodine, holmium, yttrium lithium, dye, chemical, neodymium, erbium, ruby, titanium-sapphire, diode, any  
20 harmonically oscillating laser, or any other electromagnetic radiation. Exemplary forms of heating radiation include: infrared, heating, infrared laser, radiotherapy, or any other methods of heating the lens.

Finally, exemplary forms of sound energy that can be used in an embodiment of the invention include: ultrasound, any audible and non-audible sound treatment, and any other biologically compatible sound energy.

5           The external energy used with various embodiments and methods of the present invention could be applied through either contact with the sclera or cornea, non-contact techniques, or through intraocular methods of delivery. More than one treatment may be needed to  
10 effect a suitable increase in the accommodative amplitude. When more than one modality of treatment is desirable, chemical treatment can be administered prior to, after, or simultaneously with the application of energy.

15           In an exemplary embodiment, a treatment can comprise administering a composition of one or more active agents suspended in biocompatible carrier. In another exemplary embodiment, the active agents can be administered in a solution or suspension containing  
20 ophthalmically acceptable sterile viral phage. The phage can be introduced to the lens by, for example, topical eye drop or administered systematically a pill or as an injection into either the blood stream or the

lens itself. The carrier can include, for example,  
balanced salt solution or saline. The active agents can  
include thiol transferase in an amount of 0 to 20% by  
volume, preferably 2 to 10% by volume, glutathione in an  
5 amount of 0 to 20% by volume, with a preferred range of  
2 to 10% by volume, and nicotinamide adenine  
dinucleotide phosphate (NADP) in an amount of 0-20% by  
volume, with a preferred range of 2-10% by volume. The  
balance can comprised of a biocompatible carrier. The  
10 composition can be administered in total drop volumes of  
0.1 to 2.5 ml with a referred range of 0.25 to 1 ml.

In another embodiment, thiol transferase can be  
altered to become photo reactive. Upon administering  
the composition having thiol transferase (2-10% by  
15 Vol.), glutathione (2-10% by vol.) and NADP (2-10%), a  
focused energy source such as laser can be applied to  
activate thiol transferase and the subsequent reduction  
of the disulfide bonds.